

Dynamical stabilization of runaway potentials at finite density

Alex Buchel,^{1,2,*} Junji Jia,^{2,†} and V.A. Miransky^{2,‡}

¹*Perimeter Institute for Theoretical Physics, Waterloo, Ontario N2J 2W9, Canada*

²*Department of Applied Mathematics, University of Western Ontario, London, Ontario N6A 5B7, Canada*

(Dated: January 29, 2007)

We study four dimensional non-abelian gauge theories with classical moduli. Introducing a chemical potential for a flavor charge causes moduli to become unstable and start condensing. We show that the moduli condensation in the presence of a chemical potential generates nonabelian field strength condensates. These condensates are homogeneous but non-isotropic. The end point of the condensation process is a stable homogeneous, but non-isotropic, vacuum in which both gauge and flavor symmetries and the rotational invariance are spontaneously broken. Possible applications of this phenomenon for the gauge theory/string theory correspondence and in cosmology are briefly discussed.

PACS numbers: 11.15.Ex, 11.30.Qc, 12.90.+b

It is well known [1] that in relativistic field theories introducing a chemical potential μ for a complex scalar field Φ varies its mass term from $m^2\Phi^\dagger\Phi$ to $(m^2 - \mu^2)\Phi^\dagger\Phi$. As a result, for $\mu^2 > m^2$, the field Φ condenses (the Bose-Einstein condensation). The dynamics in relativistic field theories with chemical potential for bosonic matter is rich and quite sophisticated. For example, in these theories the phenomenon of spontaneous symmetry breaking with abnormal number of Nambu-Goldstone (NG) bosons was revealed [2]. In Ref. [3], a new Higgs-like phase was described with condensates of both gauge and scalar fields which break gauge, flavor, and rotational symmetries. It is noticeable that both these phenomena have already found applications in studies of dense quark matter [4, 5].

In this Letter, we consider the role of chemical potential in non-abelian gauge theories with classical moduli. In this case, a chemical potential μ produces a negative mass squared term $-\mu^2\Phi^\dagger\Phi$, i.e., a runaway potential. Then the moduli start to condense. The question is what is the end point of this process? We will show below a possibility of an unexpected answer: the moduli condensation in the presence of a chemical potential causes the appearance of homogeneous, but non-isotropic condensates of nonabelian gauge fields in the ground states. Such condensates lead to the stable vacuum with the spontaneous breakdown of both the gauge and flavor symmetries and the rotational invariance.

The presence of moduli in nonabelian gauge theories with extended supersymmetry is a generic phenomenon.

Here, however, in order to be maximally clear, we will consider a non-supersymmetric simple model with classical moduli. Specifically, we consider the gauged $SU(2)$ σ model with a chemical potential for a complex modulus Φ in the fundamental representation of the gauge group. The chemical potential is connected with the conserved charge of the flavor hypercharge $U(1)_Y$ symmetry. The Lagrangian density is:

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}^{(a)}F^{\mu\nu(a)} + [(D_\mu - i\mu\delta_{\mu 0})\Phi]^\dagger (D_\mu - i\mu\delta_{\mu 0})\Phi, \quad (1)$$

where the hypercharge of the doublet Φ equals +1 and the $SU(2)$ gauge fields are given by $A_\mu = A_\mu^a\tau^a/2$ with τ^a being Pauli matrices. The field strength and the covariant derivative are

$$F_{\mu\nu}^{(a)} = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g\epsilon^{abc}A_\mu^b A_\nu^c, \quad D_\mu = \partial_\mu - igA_\mu. \quad (2)$$

This model is the same as that considered in Ref. [3] but with the quartic coupling constant λ and the mass m of Φ chosen to be zero. The latter makes the field Φ to be a classical modulus when $\mu = 0$.

In this model, we will demonstrate the existence of a stable vacuum with the $SU(2)_g \times U(1)_Y \times SO(3)_{\text{rot}}$ being spontaneously broken down to the $SO(2)$ rotational symmetry (here the $SU(2)_g$ is a gauge symmetry and the $SO(3)_{\text{rot}}$ is a rotational one). All our computations are classical, i.e., we neglect radiative corrections. This approximation is reliable if the chemical potential μ is much larger than the confinement scale of $SU(2)_g$, i.e., the gauge coupling g related to the scale μ is small, $g \ll 1$. Indeed, as will be shown below, in this case all the condensates are large, $\sim \mu/g$, and the dynamics corresponds to a Higgs phase at weak coupling.

A comment concerning the possibility of the one-loop Coleman-Weinberg (CW) effect [6] in this model is in

*Electronic address: abuchel@uwo.ca

†Electronic address: jjia5@uwo.ca

‡Electronic address: vmiransk@uwo.ca; On leave from Bogolyubov Institute for Theoretical Physics, 03143, Kiev, Ukraine

order. A one-loop Coleman-Weinberg (CW) potential would generate an effective quartic coupling of order g^4 . However, the important difference between the CW model (zero chemical potential) and the present one (finite chemical potential) is that while there are no tree level contributions from gauge fields in the former, the dominant effect in the present model is based on the tree level contributions of gauge bosons (vector condensates). Therefore, while in CW model the one-loop contribution of gauge fields is dominant, it is subleading in our case. In fact, one can show that a one-loop Coleman-Weinberg effective quartic potential for a complex fundamental modulus will modify our classical analysis at order $\mathcal{O}(g^2 \ln g^2)$. This correction is small at weak coupling.

We are looking for classical vacua in the σ model (1) that completely break the $SU(2)$ gauge symmetry. In this case, one can choose a unitary gauge with the vacuum expectation value of Φ taking the form

$$\Phi^T = (0, \phi_0), \quad (3)$$

where ϕ_0 is real. As in [3], we are interested in vacua with gauge field condensates breaking the $SO(3)$ rotational symmetry down to $SO(2)_{\text{rot}}$. Introducing

$$A_\mu^{(\mp)} = \frac{1}{\sqrt{2}} \left(A_\mu^{(1)} \pm i A_\mu^{(2)} \right), \quad (4)$$

such vacua are described by the condensates

$$\begin{aligned} A_3^{(+)} &= \left(A_3^{(-)} \right)^* = C \neq 0, & A_0^{(3)} &= D \neq 0, \\ A_{1,2}^{(\pm)} &= A_0^{(\pm)} = A_{1,2}^{(3)} = A_3^{(3)} = 0, \end{aligned} \quad (5)$$

for real D and complex C . We emphasize that in the unitary gauge all gauge dependent degrees of freedom are removed. Therefore here the vacuum expectation values of gauge fields are well-defined physical quantities.

The potential for the above condensates takes the form

$$V = -\mathcal{L} = -g^2 D^2 |C|^2 - \left(\mu - \frac{gD}{2} \right)^2 \phi_0^2 + \frac{g^2}{2} |C|^2 \phi_0^2, \quad (6)$$

leading to the following equations of motion

$$\left(D^2 - \frac{\phi_0^2}{2} \right) C = 0, \quad (7)$$

$$\left(2|C|^2 + \frac{\phi_0^2}{2} \right) D = \frac{\phi_0^2}{g} \mu, \quad (8)$$

$$\left[\left(\mu - \frac{gD}{2} \right)^2 - \frac{g^2}{2} |C|^2 \right] \phi_0 = 0. \quad (9)$$

The equation of motion for D in (8) represents a Gauss Law constraint for the time component of the $A^{(3)}$ field, which must be satisfied for *any* physical fluctuations. Putting D on-shell, i.e., imposing the Gauss constraint, results in the (physical) potential

$$V_{\text{phys}} = \frac{\phi_0^2 |C|^2 \left(4g^2 |C|^2 - 8\mu^2 + g^2 \phi_0^2 \right)}{2(4|C|^2 + \phi_0^2)}, \quad (10)$$

which is clearly bounded. It has the absolute minimum $V_{\text{phys}}^{(\text{min})} = -\mu^4/2g^2$ for the vacuum with

$$|C| = \frac{\mu}{\sqrt{2}g}, \quad \phi_0 = \frac{\sqrt{2}\mu}{g} \quad (11)$$

(from Gauss constraint (8), one finds $D = \frac{\mu}{g}$ in this case).

Note that there are also $SO(3)_{\text{rot}}$ -invariant solutions with

$$C = 0, \quad D = \frac{2\mu}{g}, \quad \phi_0 = \text{arbitrary}. \quad (12)$$

These $SO(3)_{\text{rot}}$ -invariant vacua (a line of moduli) with $V_{\text{phys}} = 0$ are either metastable or unstable. As will be shown below, for a sufficiently large ϕ_0 , $\phi_0 \geq 2\sqrt{2}\mu/g$, these moduli are metastable (but not unstable). For $\phi_0 < 2\sqrt{2}\mu/g$, the moduli become unstable and, as a result, the condensate C of charged vector bosons (with $Q = +1$) is dynamically generated.

At last, there is the trivial vacuum with $C = D = \Phi = 0$. It is obviously unstable.

Therefore the model (1) has the stable vacuum with nontrivial condensates (11). All gauge and flavor symmetries in this vacuum are completely broken. While the condensate ϕ_0 breaks the $SU(2)_g \times U(1)_Y$ down to the global $U(1)_{em}$ with the “electric” charge $Q = T_3 + Y/2$ [T_3 is a generator of the $SU(2)_g$], the condensate C completely breaks the $U(1)_{em}$. It also breaks the rotational $SO(3)$ down to $SO(2)_{\text{rot}}$.

It is noticeable that solution (11) describes a nonzero field strength $F_{\mu\nu}^{(a)}$ which corresponds to the presence of *non-abelian* constant “chromoelectric”-like condensates in the ground state. Choosing the vacuum with the condensate C to be real, we find

$$E_3^{(2)} = F_{03}^{(2)} = g\sqrt{2}CD = \frac{\mu^2}{g}. \quad (13)$$

We emphasize that while an abelian constant electric field in different media always leads to an instability,¹ non-abelian constant chromoelectric fields do not in many

¹ In metallic and superconducting media, such an instability is

cases. For a discussion of the stability problem for constant non-abelian fields, see the first paper in Ref. [5] and Ref. [7]. On a technical side, this difference is connected with that while a vector potential corresponding to a constant abelian electric field depends on spatial and/or time coordinates, a constant non-abelian chromoelectric field is expressed through constant vector potentials, as takes place in our case, and therefore momentum and energy are good quantum numbers in the latter.

The full spectrum of excitations in this vacuum can be explicitly computed from Lagrangian density (1) by evaluating zeroes of the determinant of the quadratic form of small fluctuations around minimum (11). One finds 10 physical states: 7 massive ones and 3 massless Nambu-Goldstone bosons associated with spontaneous breakdown of the global symmetries

$$U(1)_{em} \times SO(3)_{\text{rot}} \rightarrow SO(2)_{\text{rot}}. \quad (14)$$

The spectrum and degeneracies of massive states are given by

$$\begin{aligned} m^2 &= 2\mu^2 & [\times 2], \\ m^2 &= 5\mu^2 & [\times 2], \\ m^2 &= 4\mu^2 & [\times 1], \\ m^2 &= \gamma_-^2 \mu^2 & [\times 1], \\ m^2 &= \gamma_+^2 \mu^2 & [\times 1], \end{aligned} \quad (15)$$

where

$$\gamma_{\pm}^2 = \frac{5 \pm \sqrt{13}}{2}. \quad (16)$$

Note that since all the gauge and flavor symmetries and the rotational $SO(3)$ are broken in this vacuum, it describes an anisotropic superconducting medium. Its dynamics should be very rich [8].²

We also calculated the spectrum of excitations in the $SO(3)$ invariant vacua (12). Their spectrum includes a single massless ϕ_0 -modulus excitation and 9 massive modes assigned to three $SO(3)_{\text{rot}}$ triplets (vector modes). Their masses are $m_0 = g\phi_0$ and $m_{\pm} = (g\phi_0 \pm 2\sqrt{2}\mu)$. While the mass m_0 is connected with the neutral $A^{(3)}$ vector boson ($Q = 0$), $m_{\pm}$ are the masses of charged $A^{(\mp)}$ vector bosons ($Q = \mp 1$). For a large

scalar condensate $\phi_0 > 2\sqrt{2}\mu/g$, all the masses are positive and therefore the vacua are not unstable, although metastable: while their (free) energy density V_{phys} is zero, $V_{\text{phys}}^{(\text{min})}$ in the ground state (11) is negative. On the other hand, at the values of ϕ_0 less than $2\sqrt{2}\mu/g$, the mass m_- is negative and therefore the process of a crossover of particle-antiparticle levels takes place. The latter is a signature of the Bose-Einstein instability: at these values of ϕ_0 , the condensate C of charged $A^{(+)}$ vector bosons is dynamically generated.

Thus, in this simple but nontrivial model, we showed that unstable directions in nonabelian gauge theories induced by a chemical potential for the matter density can be stabilized by the generation of nonabelian gauge field strength condensates. Such condensates are homogeneous but non-isotropic. It would be interesting to extend our analysis to more general gauge groups and matter representations and to understand the general conditions under which dynamical stabilization of the runaway directions along the lines proposed in this paper can occur [9].

We would like to conclude with outlining three specific research directions. First, consider the $\mathcal{N} = 4$ supersymmetric $SU(2)$ Yang-Mills (SYM) theory in the presence of the $U(1)$ R-charge chemical potential. This model is similar to the gauged σ model (1). In fact, on the $\mathcal{N} = 4$ moduli space, this would be precisely σ model (1) but with three copies of the complex adjoint scalar fields instead of a single fundamental Higgs Φ . It is straightforward to repeat our analysis for this SYM and demonstrate that moduli condensation here also induces non-abelian field strength condensates.³ Following the gauge theory/string theory correspondence [10], a stable non-isotropic vacuum in $\mathcal{N} = 4$ SYM in the presence of a chemical potential (if it exists) would imply a new ground state for a configuration of rotating D3-branes with non-isotropic world-volume.

Secondly, while in this Letter we discussed four-dimensional gauge theories in which the spontaneous rotational symmetry breakdown is generated due to finite density of matter fields, four-dimensional gravitational effects were completely neglected. Coupling of a higher dimensional gauge theory with finite matter density to gravity could produce stable vacua where higher dimensional rotational invariance is spontaneously broken down to the effective low-energy $SO(3)$ rotational invariance of

classical in its origin. In semiconductors and insulators, this instability is manifested in creation of electron-hole pairs through a quantum tunneling process.

² As has been shown recently in Ref. [9], the solution (11) yields the global vacuum in this model.

³ The pattern of the gauge and rotational symmetries breaking appears to be more complicated in this case. This model will be discussed elsewhere.

our Universe.

Finally, the phenomena described in this Letter occurs at high energy, i.e., for weak coupling of an asymptotically free gauge theory. Thus it might have important observable signatures in the cosmological evolution of our Universe.

AB would like to thank Jerome Gauntlett, Maxim Pospelov and Alfred Shapere for valuable discussions. AB would like to thank the Aspen Center for Physics for

hospitality during the course of this work. AB research at Perimeter Institute is supported in part by the Government of Canada through NSERC and by the Province of Ontario through MEDT. AB gratefully acknowledges further support by an NSERC Discovery grant. VAM thanks Igor Shovkovy for useful discussions. The work of JJ and VAM was supported by the Natural Sciences and Engineering Research Council of Canada.

-
- [1] J. I. Kapusta, Phys. Rev. D **24**, 426 (1981); H. E. Haber and H. A. Weldon, Phys. Rev. D **25**, 502 (1982).
 - [2] V. A. Miransky and I. A. Shovkovy, Phys. Rev. Lett. **88**, 111601 (2002), hep-ph/0108178; T. Schafer, D. T. Son, M. A. Stephanov, D. Toublan and J. J. M. Verbaarschot, Phys. Lett. B **522**, 67 (2001), hep-ph/0108210.
 - [3] V. P. Gusynin, V. A. Miransky and I. A. Shovkovy, Phys. Lett. B **581**, 82 (2004), hep-ph/0311025; Mod. Phys. Lett. A **19**, 1341 (2004), hep-ph/0406219.
 - [4] D. Blaschke, D. Ebert, K. G. Klimenko, M. K. Volkov and V. L. Yudichev, Phys. Rev. D **70**, 014006 (2004), hep-ph/0403151; D. Ebert, K. G. Klimenko and V. L. Yudichev, Phys. Rev. D **72**, 056007 (2005), hep-ph/0504218; *ibid* D **75**, 025024 (2007), hep-ph/0608304; L. He, M. Jin and P. Zhuang, hep-ph/0504148; A. H. Rezaeian and H. J. Pirner, Nucl. Phys. A **779**, 197 (2006), nucl-th/0606043.
 - [5] E. V. Gorbar, M. Hashimoto and V. A. Miransky, Phys. Lett. B **632**, 305 (2006), hep-ph/0507303; E. V. Gorbar, M. Hashimoto, V. A. Miransky and I. A. Shovkovy, Phys. Rev. D **73**, 111502(R) (2006), hep-ph/0602251; K. Fukushima, Phys. Rev. D **73**, 094016 (2006), hep-ph/0603216; O. Kiriyama, D. H. Rischke and I. A. Shovkovy, Phys. Lett. B **643**, 331 (2006), hep-ph/0606030.
 - [6] S. R. Coleman and E. Weinberg, Phys. Rev. D **7**, 1888 (1973).
 - [7] L. S. Brown and W. I. Weisberger, Nucl. Phys. B **157**, 285 (1979) [Erratum-*ibid.* B **172**, 544 (1980)].
 - [8] Based on the analysis in E. V. Gorbar, J. Jia and V. A. Miransky, Phys. Rev. D **73**, 045001 (2006), hep-ph/0512203, we expect that there exist different types of vortices in this model.
 - [9] A. Buchel, J. Jia and V. A. Miransky, hep-th/0702066.
 - [10] J. M. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998), Int. J. Theor. Phys. **38**, 1113 (1999), hep-th/9711200; O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, Phys. Rept. **323**, 183 (2000), hep-th/9905111.